

**TRANSFER IN GALOIS COHOMOLOGY COMMUTES WITH
TRANSFER IN THE MILNOR RING**

Jack SHAPIRO

Department of Mathematics, Washington University, St. Louis, MO 63130, USA

Communicated by H. Bass

Received April 1979

The purpose of this paper is to study the relation between transfer maps defined in the theory of quadratic forms, the algebraic K -theory of Milnor and Galois cohomology, respectively. The results should be viewed as an addendum to [8] and to an unpublished letter of John Tate.

For a field F , $\text{ch}(F) \neq 2$, we have an associated ring, $\hat{W}(F)$, called the Witt–Grothendieck ring of regular quadratic forms over F [4]. If E is a finite extension of F then the trace linear functional induces a transfer map, $T: \hat{W}(E) \rightarrow \hat{W}(F)$, which is an additive group homomorphism [4, Chapter 7]. Now consider $K_*(F)$, the Milnor ring of F . For any finite separable extension $E = F(\alpha)$ of F we have an induced map $N_\alpha: K_*(E) \rightarrow K_*(F)$ defined by Bass and Tate in [1]. It is as yet unknown whether N_α is independent of the choice of primitive element α ¹. If $k_*(F) = K_*(F)/2K_*(F)$, the reduced Milnor ring, then we will show that if E is Galois over F of odd degree then $N_\alpha: k_*(E) \rightarrow k_*(F)$ is indeed independent of α . In fact we will show that the restriction of N_α to $k_2(E)$ is the canonical transfer map (mod 2) defined by Milnor in [7].

Let $H^*(F) = H^*(F; \mathbb{Z}/2\mathbb{Z})$ be the Galois cohomology ring of the absolute Galois group of F with coefficients in $\mathbb{Z}/2\mathbb{Z}$. If $\text{Gal}(\bar{F}/F)$ is the Galois group of a separable closure \bar{F} over F then in terms of group cohomology

$$H^*(\text{Gal}(\bar{F}/F)) = \varinjlim_{i \in \Lambda} H^*(\text{Gal}(F_i/F))$$

where $\{F_i\}_{i \in \Lambda}$ is the set of all finite Galois extensions of F contained in \bar{F} ([9]). In this case corestriction gives us a canonical transfer map, $\text{cores}: H^*(E) \rightarrow H^*(F)$ whenever E is a finite extension of F .

There are two main results in this paper. The first generalizes a result of Tate which appears in a letter to S. Rosset. For any field F we have a ring homomorphism $h_F: k_*(F) \rightarrow H^*(F)$ [6; Lemma 6.1]. If E is a Galois extension of F of odd degree, $E = F(\alpha)$, let $N = N_\alpha$ be the canonical transfer map.

¹ K. Kato has recently proved the independence of the choice of α (unpublished manuscript).

Theorem 1. For any $n \geq 0$

$$\begin{array}{ccc} k_n(E) & \xrightarrow{h_E} & H^n(E) \\ \downarrow N & & \downarrow \text{cores} \\ k_n(F) & \xrightarrow{h_F} & H^n(F) \end{array}$$

is a commutative diagram of abelian groups.

If we assume that for any Galois extension $E = F(\alpha)$ the transfer map N_α is independent of α and is also functorial with respect to a tower of extensions then we could drop the 'odd degree' assumption in Theorem 1. Tate proved a slightly different result which covers the case $n = 2$ with N being the above cited transfer map of Milnor. The cases $n = 0, 1$ are trivial.

The second theorem is a corollary to the main theorem of [8]. Let w_i , $i = 0, 1, 2, 3, \dots$, be the Stiefel–Whitney invariants defined by Delzant–Milnor and let us view them as maps $w_n: \hat{W}(F) \rightarrow H^n(F)$. $\hat{I}(F)$ will denote the augmentation ideal, i.e. the kernel of the dimension homomorphism $\dim: \hat{W}(F) \rightarrow \mathbb{Z}$. If $t = 2^{n-1}$ then $w_t: \hat{I}^n(F) \rightarrow H^t(F)$ is a group homomorphism.

Theorem 2.

$$\begin{array}{ccc} \hat{I}^n(E) & \xrightarrow{w_t} & H^t(E) \\ \downarrow T & & \downarrow \text{cores} \\ \hat{I}^n(F) & \xrightarrow{w_t} & H^t(F) \end{array}$$

is a commutative diagram of abelian groups. $\hat{I}^n(F)$ is the n -th power ideal of $\hat{I}(F)$ and T is the transfer map restricted to $\hat{I}^n(E) \leq \hat{W}(E)$.

1.

For any field F , $\text{ch}(F) \neq p$, Tate has constructed an associated field F' which will play an important role in what follows. The construction is introduced in [10] but in spirit it goes back to the methods used in [1, Section 5]. For this paper we are basically interested in the case $p = 2$.

Consider the set of all subextensions of a separable closure, \bar{F} , of F which (1) contain F and (2) can be realized as a set-theoretic union of finite prime-to- p extensions of F . A Zorn's lemma argument implies that this set contains a maximal element. Call it F' . This field F' has some very interesting and useful properties. These properties are asserted in Tate's letter. Since the letter remains unpublished

we will *sketch* the proofs below. I would like to thank Jack Sonn and Roger Ware for conversations which helped me better understand the nature of F' .

Property 1. *If E is any finite extension of F contained in F' then the degree $[E:F]$ is prime to p .*

Proof. $E = F(\alpha_1, \dots, \alpha_r)$ where $\alpha_i \in F'$. Hence their irreducible polynomials over F must be of degree prime to p .

Property 2. *If E is any finite extension of F' contained in \bar{F} then $[E:F']$ is equal to p^a for some nonnegative integer a .*

Proof. This will follow if we show that for any $\alpha \in E$ the irreducible polynomial of α over F' , $\text{Irr}(\alpha, F')$, has as its degree a power of p . All the coefficients of $\text{Irr}(\alpha, F')$ lie in finite prime-to- p extensions of F . If the degree of α over F' is also prime-to- p a straightforward argument shows that $F'(\alpha)$, a proper extension of F' , contradicts the maximality of F' . Assume that the degree of α over F' is $p^a n$, $a \geq 1$, $n \neq 0$ and $(p, n) = 1$. Let E_N be the normal closure of E in \bar{F} . It is a Galois extension of F' whose degree over F' is divisible by $p^a n$. By the above assumptions, a Sylow p -subgroup, $S(p)$, of $\text{Gal}(E_N/F')$ is a nontrivial proper subgroup. Therefore the fixed subfield, $E_N^{S(p)}$, would be a nontrivial prime-to- p extension of F' which we have already seen cannot happen. It follows that $n = 1$ and we are done.

Property 3. *Let E be a finite extension of F' contained in \bar{F} , $[E:F'] = p^a$. Then there exists a tower of fields*

$$F' = E_1 \subset E_2 \subset \dots \subset E_a = E$$

such that $[E_i : E_{i-1}] = p$.

Proof (by induction on a). To establish the property we need only find a subfield K of E whose degree over F' is p^{a-1} . $G = \text{Gal}(\bar{F}/F')$ is a pro- p -group since by Property 2 all finite extensions of F' contained in \bar{F} are p -power extensions. Galois theory implies that E is the fixed subfield of a subgroup H of G where $[G:H] = p^a$. We will find a subgroup H_1 , $H \leq H_1 \leq G$ so that $[G:H_1] = p^{a-1}$. Letting $K = \bar{F}^{H_1}$ we will get the desired subfield K .

H is an open subgroup of G (finite index). By the class equation it also follows that H has only a finite number of conjugates in G . Let $H' = \bigcap_{x \in G} x^{-1} H x$, then H' is an open normal subgroup of G containing H . G/H' is a finite p -group containing H/H' as a subgroup. By the Sylow theorems we can find H_1 , normal in G , with $H' \leq H_1 \leq H$ and $[G:H_1] = p^{a-1}$. This then completes the proof.

Let us note that from the above proof it follows that if we assume E/F is Galois we can construct the E_i so that each is Galois over E_{i-1} .

Property 4. *Let $p=2$. Then the restriction map $\text{res} : H^*(F) \rightarrow H^*(F')$ is injective.*

Proof. Since coefficients are in $\mathbb{Z}/2\mathbb{Z}$ it is well known that for finite groups $G' \leq G$, $\text{res} : H^*(G) \rightarrow H^*(G')$ is injective if the index of G' in G is odd (prime to 2). By Property 1 every finite extension of F in F' is odd degree. Using the definition of Galois cohomology as a direct limit of finite group cohomology, injectivity for $\text{res} : H^*(F) \rightarrow H^*(F')$ follows.

2.

Following the notation of [6] $K_n(F)$ will denote the Milnor K -group of n -fold symbols over F . $K_n(F)$ is additively generated by symbols $l(a_1) \cdots l(a_n)$ where $a_i \in F^*$ is a unit in F . The quotient group $K_n(F)/2K_n(F)$ will be denoted by $k_n(F)$. $k_*(F)$ is the reduced Milnor ring and as a group is isomorphic to $\coprod_{n \geq 0} k_n(F)$. By $k_n(F)$ we mean the $\mathbb{Z}/2\mathbb{Z}$ algebra consisting of all formal series $\gamma_0 + \gamma_1 + \gamma_2 + \cdots$ with $\gamma_i \in k_i(F)$.

Suppose E is a finite separable extension of F with primitive element α , i.e. $E = F(\alpha)$. Bass and Tate define a transfer map $N_\alpha : K_n(E) \rightarrow K_n(F)$. By reducing mod $2K_n(E)$ we have $N_\alpha : k_n(E) \rightarrow k_n(F)$. The question is whether N_α is independent of α and if so is it also functorial. By functorial we mean that if $E \supset K \supset F$ is a tower of separable extensions with $K = F(\alpha)$, $E = K(\beta)$ and $E = F(\gamma)$ then $N_\gamma = N_\alpha \circ N_\beta$. A partial solution is suggested in [1] but to my knowledge the question still remains open. What we will now show is that we can answer the first question in the affirmative if E is Galois over F of odd degree.

Proposition 1. *Let E be a finite Galois extension of F of odd degree and suppose $E = F(\alpha) = F(\beta)$. Then $N_\alpha = N_\beta : k_*(E) \rightarrow k_*(F)$.*

Proof. The proof is essentially an application of the partial solution suggested by Bass and Tate and so we will refer the reader to [1, pp. 39–40]. For the field L in [1] substitute the field F' constructed in the previous section. Since F' is a union of odd-degree extensions the restriction map $j : k_*(F) \rightarrow k_*(F')$ is injective by [1, p. 40]. By the standard theorems of Galois theory (see for example [5]) $[F'(\alpha_i) : F']$ divides $[F(\alpha) : F]$ for all i . Since $[F(\alpha) : F]$ is odd and since F' has only 2-power degree extensions it follows that for all i , $F'(\alpha_i) = F'$. The general case now follows from the case of degree 1 extensions where $N_\alpha \equiv \text{identity}$.

3.

For a field F , $\text{ch}(F) \neq 2$, with separable closure \bar{F} let G be the absolute Galois group of F , i.e. $G = \text{Gal}(\bar{F}/F)$. By [6, Section 6] we have a group homomorphism

$\delta: F^*/(F^*)^2 \rightarrow H^1(G)$, where F^* is the multiplicative group of units in the field F . By viewing $k_1(F)$ as $F^*/(F^*)^2$ and by using the cup product in cohomology we get an induced ring homomorphism $h_F: k_*(F) \rightarrow H^*(G)$. For each $n \geq 0$, $h_F: k_n(F) \rightarrow H^n(G)$ is a group homomorphism. Following the usual notation we will denote $H^*(G)$ by $H^*(F)$ (F completely determines the isomorphism class of G). There are many examples for which h_F is known to be bijective, e.g. finite, local, global or real closed fields [6, Lemma 6.2].

As we noted previously (Proposition 1), if E is a Galois extension of F of odd degree then we have a canonical transfer map $N: k_n(E) \rightarrow k_n(F)$ for any $n \geq 0$.

Theorem 1. *For E and F as above we have the following commutative diagram of groups and group homomorphisms:*

$$\begin{array}{ccc} k_n(E) & \xrightarrow{h_E} & H^n(E) \\ \downarrow N & & \downarrow \text{cores} \\ k_n(F) & \xrightarrow{h_F} & H^n(F) \end{array} \quad (1)$$

where *cores* is the corestriction map in cohomology.

Remark. A stronger result for the case $n=2$ is proved by Tate in [10]. Our proof will be an application of Tate's methods to our particular situation. The idea is to create a cube whose vertices are groups and whose edges are group homomorphisms in such a way that diagram (1) is one of the faces. By a sequence of lemmas we will show that all the other faces commute. This in turn will imply that (1) is a commutative square.

In proving the lemmas we will only assume that E is a finite separable extension of F . For the theorem itself we will need both the assumptions that E is Galois and of odd degree over F . We have $E = F(\alpha)$ for some $\alpha \in E$ and a possibly noncanonical transfer map $N = N_\alpha: k_n(E) \rightarrow k_n(F)$ (see Section 2). Let L be an extension of F contained in \bar{F} . Note that although L is arbitrary we will eventually apply the lemmas to the case where $L = F'$, the special extension of F constructed in Section 1. Let \tilde{L} be defined as $E \otimes_F L$. \tilde{L} can be realized as a direct product of fields, $\tilde{L} \cong \prod_{i=1}^r L_i$, where each L_i is a finite separable extension of L . If we assume that E is Galois over F then each L_i is Galois over L and furthermore $[L_i: L]$ divides $[E: F]$. If we let $p(t) = \text{Irr}(\alpha, F)$ then in $L[t]$ $p(t)$ will decompose into a product $p_1(t) \cdots p_r(t)$ where each $p_i(t)$ is a monic irreducible polynomial. We can take as L_i the quotient $L[t]/(p_i(t))$ where $(p_i(t))$ represents the principal ideal generated by $p_i(t)$. In general \tilde{L} is not a field (unless $p(t)$ is irreducible in $L[t]$) and so we need to make some definitions. For $n \geq 0$ we will define $k_n(\tilde{L})$ as a direct product $\prod_{i=1}^r k_n(L_i)$ and $H^n(\tilde{L})$ as $\prod_{i=1}^r H^n(L_i)$. When we refer to a restriction, corestriction or transfer map we will mean the appropriate sum or direct product maps. For example $N: k_n(\tilde{L}) \rightarrow k_n(L)$ is

the sum in $k_n(L)$ of the maps $N_{\alpha_i} : k_n(L_i) \rightarrow k_n(L)$ where α_i is a root of $p_i(t)$ in L_i . By $\text{res} : k_n(E) \rightarrow k_n(\tilde{L})$ we mean the direct product map of the $\text{res}_i : k_n(E) \rightarrow k_n(L_i)$.

Lemma 1.

$$\begin{array}{ccc} k_n(E) & \xrightarrow{\text{res}} & k_n(\tilde{L}) \\ \downarrow N & & \downarrow N \\ k_n(F) & \xrightarrow{\text{res}} & k_n(L) \end{array}$$

is a commutative diagram of groups and group homomorphisms.

Proof. The diagram is one portion of the commutative exact sequence diagram of [1, Section 5.8].

Lemma 2. Let $\text{res} : H^n(E) \rightarrow H^n(\tilde{L})$ be the direct product map $\prod_{i=1}^r \text{res}_i$ where $\text{res}_i : H^n(E) \rightarrow H^n(L_i)$. Let $\text{cores} : H^n(\tilde{L}) \rightarrow H^n(L)$ be the sum in $H^n(L)$ of the maps $\text{cores}_i : H^n(L_i) \rightarrow H^n(L)$. Then the following diagram of groups and group homomorphisms is commutative:

$$\begin{array}{ccc} H^n(E) & \xrightarrow{\text{res}} & H^n(\tilde{L}) \\ \downarrow \text{cores} & & \downarrow \text{cores} \\ H^n(F) & \xrightarrow{\text{res}} & H^n(L) \end{array}$$

Proof. Let $G_F = \text{Gal}(\bar{F}/F)$ and let us use the notation of group cohomology to write $H^n(F) = H^n(G_F)$. Let $\alpha = \alpha_1, \alpha_2, \dots, \alpha_s$ be the roots of $p(t) = \text{Irr}(\alpha, F)$ in \bar{F} . In $L[t]$ $p(t) = p_1(t) \cdots p_r(t)$. Choose the ordering so that $\alpha = \alpha_1$ is a root of $p_1(t)$. Then the composite EL of E and L in \bar{F} is isomorphic to $L_1 \cong L[t]/(p_1(t))$. By separability all the α_i are distinct.

Partition the roots into r disjoint subsets $S_i = \{\alpha_{i1}, \dots, \alpha_{is_i}\}$, $1 \leq i \leq r$, where S_i is the set of roots of $p_i(t)$ in \bar{F} . In particular we have $\sum_{i=1}^r s_i = s$. We may order S_i so that $\alpha_{i1} = \alpha$. For each $1 \leq i \leq r$ there exists an embedding η_i of E in \bar{F} over F such that $\eta_i(\alpha_{i1}) = \alpha_{i1}$. Let $\eta_i(E)$ be denoted by E_i . Then $E_i = F(\alpha_{i1})$ and L_i is isomorphic to the composite $E_i L$.

For each i , $1 \leq i \leq r$, we have the following diagram of fields:

$$\begin{array}{ccccc} E & \xrightarrow{\eta_i} & E_i & \xrightarrow{\quad} & E_i L \cong L_i \\ & \searrow & \nearrow & & \nearrow \\ & & F & \xrightarrow{\quad} & L \end{array}$$

and a corresponding diagram for cohomology

$$\begin{array}{ccccc}
 H^n(G_E) & \xrightarrow{\eta_i^*} & H^n(G_{E_i}) & \xrightarrow{\text{res}_i} & H^n(G_{L_i}) \\
 \searrow \text{cores} & & & & \swarrow \text{cores}_i \\
 & & H^n(G_F) & \xrightarrow{\text{res}} & H^n(G_L)
 \end{array}$$

η_i induces a map from $G_{E_i} \rightarrow G_E$ and η_i^* is the induced map on cohomology. To prove the lemma we need to show that

$$\text{res} \circ \text{cores} = \sum_{i=1}^r \text{cores}_i \circ \text{res}_i \circ \eta_i^*. \quad (2)$$

Let us recall the definition of corestriction given in [11]. Given a profinite group G and a subgroup G' of finite index choose a set of representatives for the right cosets of G' in G (a right transversal). If R is the right transversal chosen then $G = \bigcup_{r \in R} G'r$. On the cochain level define

$$\text{cores}(f)(X_1, \dots, X_n) = \sum_{r \in R} f(r^{(0)}X_1 r^{(1)-1}, \dots, r^{(n-1)}X_n r^{(n)-1}),$$

where f is a G' cochain and $(X_1, \dots, X_n) \in G^n$. The $r^{(i)}$ are determined inductively by $r^{(0)} = r$ and for $1 \leq i \leq n$ $r^{(i)} \in G'r^{(i-1)}X_i$, $r^{(i)} \in R$. As noted in [11] this defines a map on cohomology independent of the choice of R . As it will turn out formula (2) is valid even on the cochain level.

For each i, j , $1 \leq i \leq r$, $1 \leq j \leq s_i$, let $\tau_{ij} \in G_L$ be chosen so that $\tau_{ij}(\alpha_{ij}) = \alpha_{i1}$. For fixed i the set $\{\tau_{ij}\}_{j=1}^{s_i}$ gives a right transversal for G_{L_i} in G_L . It follows that $\{\eta_i^{-1}\tau_{ij}\}_{i=1}^r \}_{j=1}^{s_i}$ provides a right transversal for G_E in G_F . Let f be an n -cochain representing an element in $H^n(G_E)$. For $(X_1, \dots, X_n) \in G_L^n$ we have

$$\begin{aligned}
 \sum_{i=1}^r \text{cores}_i \circ \text{res}_i \circ \eta_i^*(f)(X_1, \dots, X_n) &= \\
 &= \sum_{i=1}^r \sum_{j=1}^{s_i} \text{res}_i \circ \eta_i^*(f)(\tau_{ij}^{(0)}X_1 \tau_{ij}^{(1)-1}, \dots, \tau_{ij}^{(n-1)}X_n \tau_{ij}^{(n)-1}) \\
 &= \sum_{i=1}^r \sum_{j=1}^{s_i} f(\eta_i^{-1}\tau_{ij}^{(0)}X_1(\eta_i^{-1}\tau_{ij}^{(1)})^{-1}, \dots, \eta_i^{-1}\tau_{ij}^{(n-1)}X_n(\eta_i^{-1}\tau_{ij}^{(n)})^{-1}) \\
 &= \text{res} \circ \text{cores}(f)(X_1, \dots, X_n)
 \end{aligned}$$

which is what we needed.

Lemma 3. *The following is a commutative diagram of rings and ring homomorphisms:*

$$\begin{array}{ccc}
 k_*(L) & \xrightarrow{h_L} & H^*(L) \\
 \uparrow \text{res} & & \uparrow \text{res} \\
 k_*(F) & \xrightarrow{h_F} & H^*(F)
 \end{array}$$

In particular if we put any $n > 0$ in place of $*$ we get a commutative diagram of abelian groups.

Proof. Since $k_*(F)$ is generated as a ring by $k_1(F)$ it is enough to check the diagram for $n = 1$. $h_F: k_1(F) \cong F^*/(F^*)^2 \rightarrow H^1(F) \cong \text{Hom}(G_F; \mathbb{Z}/2\mathbb{Z})$ is defined for any $\alpha \in F^*$ and $\sigma \in G_F$ by $h_F(\alpha)(\sigma) = \sigma(\sqrt{\alpha})(\sqrt{\alpha})^{-1}$ where $\sqrt{\alpha}$ is any root of $x^2 - \alpha$ in F . A straightforward check shows that the diagram commutes for $n = 1$ and the rest follows as indicated.

Apply Lemma 3 to all the extensions L_i over E for $i = 1, \dots, r$. Then take

$$\text{res} = \prod_{i=1}^r \text{res}_i : H^n(E) \rightarrow \prod_{i=1}^r H^n(L_i)$$

and

$$h_L = \prod_{i=1}^r h_{L_i} : \prod_{i=1}^r k_n(L_i) \rightarrow \prod_{i=1}^r H^n(L_i).$$

We then get the following additional lemma:

Lemma 4. For any $n > 0$

$$\begin{array}{ccc}
 k_n(\tilde{L}) & \xrightarrow{h_{\tilde{L}}} & H^n(\tilde{L}) \\
 \uparrow \text{res} & & \uparrow \text{res} \\
 k_n(E) & \xrightarrow{h_E} & H^n(E)
 \end{array}$$

is a commutative diagram.

Proof of Theorem 1. We now restrict our attention to the case where E/F is Galois of odd degree. We choose as our extension L the extension F' constructed in Section 1. We can construct a cube whose vertices are groups and whose edges are group homomorphisms in such a way that diagram (1) of Theorem 1 is one of the faces. The four diagrams of Lemmas 1–4 will provide four other faces. The sixth face is

$$\begin{array}{ccc}
 \prod_{i=1}^r k_n(L_i) & \xrightarrow{\prod_{i=1}^r h_{L_i}} & \prod_{i=1}^r H^n(L_i) \\
 \downarrow N & & \downarrow \text{cores} \\
 k_n(L) & \xrightarrow{h_L} & H^n(L)
 \end{array} \tag{3}$$

where N is the sum in $k_n(L)$ of the transfer maps N_i and cores is the sum in $H^n(L)$ of cores_i . Since $L = F'$ is a union of odd degree extensions of F it follows that $\text{res} : H^n(F) \rightarrow H^n(L)$ is injective (Property 4). The commutativity of diagram (1) will therefore follow from the commutativity of (3) together with the results of Lemma 1–4. The extension E/F being Galois of odd degree it therefore follows that L_i/L is Galois of odd degree. Property 2 of Section 1 says that any nontrivial finite extension of $L = F'$ must be of 2-power order. Therefore $[L_i : L] = 1$ for all $1 \leq i \leq r$. The commutativity of (3) is then immediate.

Remarks. If we assume that for any finite separable extension $F(\alpha)/F$ the map N_α is independent of α and is also functorial for a tower of extensions, then Theorem 1 is valid for any finite separable extension. The observations are as follows. We have already seen that the commutativity of diagram (1) follows from that of diagram (3). By Property 2 if $L = F'$ then each L_i must be a 2-power separable extension of L . By functoriality and Property 3 we need only consider the case where each L_i is a quadratic extension of L . We are therefore reduced to proving the commutativity of

$$\begin{array}{ccc} k_n(E) & \xrightarrow{h_E} & H^n(E) \\ \downarrow N & & \downarrow \text{cores} \\ k_n(F) & \xrightarrow{h_F} & H^n(F) \end{array} \quad (4)$$

where $[E : F] = 2$. By [1, Corollary 5.3], $k_*(E)$ is generated as a $k_*(F)$ -module by $k_1(E)$. We also know that N is a $k_*(F)$ -module homomorphism ($\text{res} : k_*(F) \rightarrow k_*(E)$ makes $k_*(E)$ into a $k_*(F)$ -module). Finally $N : k_1(E) \cong E^*/(E^*)^2 \rightarrow k_1(F) \cong F^*/(F^*)^2$ is the classical norm map. Putting these facts together implies that (4) is a commutative diagram. The problem that remains is to define a canonical and functorial transfer map for any finite separable extension E over F .

4.

We now turn our attention to the second theorem described in the introduction. Let $\hat{I}(F) = \ker\{\dim : \hat{W}(F) \rightarrow Z\}$, the augmentation ideal, and let its n -th power be denoted by $\hat{I}^n(F)$. Delzant [2] and Milnor [6] have generalized the classical invariants dimension, discriminant and the Hasse invariant to give an infinite sequence of invariants w_n , $n = 0, 1, 2, \dots$. These Stiefel–Whitney invariants define maps (not necessarily homomorphisms) from $\hat{W}(F)$ to $k_n(F)$ or $H^n(F)$ depending on whether one follows Milnor's or Delzant's definition. The connection is that for $q \in \hat{W}(F)$, $w_n(q) \in H^n(F)$ of Delzant equals $h_F(w_n(q))$, where $w_n(q) \in k_n(F)$ is as defined by Milnor. Recall now the definition of $k_n(F)$ as given in Section 2. Define $H^n(F)$ analogously. The w_i induce a homomorphism $w : \hat{W}(F) \rightarrow k_n(F)^*$ (or $H^n(F)^*$).

$w = w_0 + w_1 + w_2 + \cdots$ is called the total Stiefel–Whitney class. The homomorphism is from the additive group $\hat{W}(F)$ to the multiplicative group of units in $k_\pi(F)$ (or $H^\pi(F)$). From now on when we use the w_i we will view them as maps from $\hat{W}(F)$ to $H^i(F)$. Note that although w_i is not a homomorphism, if $t = 2^{n-1}$, $n \geq 1$, then by [6, Lemma 3.2], $w_t: \hat{I}^n(F) \rightarrow H^t(F)$ is a group homomorphism.

Theorem 2. *If $t = 2^{n-1}$ where $n \geq 1$, then*

$$\begin{array}{ccc} \hat{I}^n(E) & \xrightarrow{w_t} & H^t(E) \\ \downarrow \tau & & \downarrow \text{cores} \\ \hat{I}^n(F) & \xrightarrow{w_t} & H^t(F) \end{array}$$

is a commutative diagram of groups and group homomorphisms.

The proof of Theorem 2 will follow as a corollary to the main theorem of [8]. Besides corestriction there exists a multiplicative transfer map $\mathcal{N}: H^\pi(E) \rightarrow H^\pi(F)$. \mathcal{N} is a multiplicative group homomorphism from the group of units of $H^\pi(E)$ to the group of units of $H^\pi(F)$ [8]. The map \mathcal{N} is defined as a direct limit of $\mathcal{N}_i: H^\pi(\text{Gal}(E_i/E)) \rightarrow H^\pi(\text{Gal}(F_i/F))$ where $\{E_i\}_{i \in \Lambda}$ is the set of all finite Galois extensions of E contained in F . Recall that $H^\pi(E) \cong \varinjlim_{i \in \Lambda} H^\pi(\text{Gal}(E_i/E))$. For each $i \in \Lambda$, $\text{Gal}(E_i/E)$ is a subgroup of the finite group $\text{Gal}(E_i/F)$, of index $l = [E:F]$. For a finite group the definition of \mathcal{N}_i goes back to Evens [3].

Theorem (main theorem of [8]). *Let E be a finite Galois extension of F . If $[E:F]$ is odd then the diagram*

$$\begin{array}{ccc} \hat{W}(E) & \xrightarrow{w} & H^\pi(E) \\ \downarrow \tau & & \downarrow \mathcal{N} \\ \hat{W}(F) & \xrightarrow{w} & H^\pi(F) \end{array} \quad (5)$$

is commutative. If $[E:F]$ is even, then the diagram commutes modulo the kernel of the restriction map, $\text{res}: H^\pi(F) \rightarrow H^\pi(E)$.

We now introduce a lemma which will relate \mathcal{N} to the corestriction homomorphism.

Lemma. *Let G be a profinite group and let G' be a subgroup of G of finite index. Suppose $\gamma \in H^\pi(G)$ is of the form $1 + \alpha + \alpha_1 + \alpha_2 + \cdots$ where $\alpha \in H^\pi(G')$ and for $i = 1, 2, 3, \dots$, $\alpha_i \in H^i(G')$ with $i > n$. Then*

$$\mathcal{A}(\gamma) = 1 + \text{cores}(\gamma) + \beta_1 + \beta_2 + \dots$$

where $\beta_j \in H^j(G)$ and $j > n$ for $j = 1, 2, \dots$

Proof.

Since \mathcal{A} is defined as a direct limit, as noted, it is enough to prove the lemma for the case where G is a finite group. We must also note of course that for a profinite group corestriction can be defined as a direct limit of the corestriction maps on each finite level. For the case of a finite group the lemma becomes a slight generalization of Theorem 1 in [3].

Let $1 = \tau_1, \dots, \tau_l$ be a left transversal of G' in G . This set induces a group homomorphism $\Phi: G \rightarrow S_l \tilde{X}(G')^l$, the semi-direct product of the symmetric group S_l and $(G')^l$. The action of S_l is by permuting the entries of $(G')^l$ [3, p. 55]. Let

$$W = \dots \rightarrow W_i \rightarrow W_{i-1} \rightarrow \dots \rightarrow W_0 \xrightarrow{\varepsilon} Z \rightarrow 0$$

be an S_l -projective resolution of Z and let

$$X = \dots \rightarrow X_i \rightarrow X_{i-1} \rightarrow \dots \rightarrow X_0 \xrightarrow{\mu} Z \rightarrow 0$$

be a G' -projective resolution of Z . Then $W \otimes X^l$ (tensor product of resolutions) is an $S_l \tilde{X}(G')^l$ projective resolution of Z and by the map Φ also a G projective resolution of Z .

Define $\eta: Z \rightarrow Z/2Z$ by $\eta(1) = 1 \pmod{2}$ then $\mu' = \eta \circ \mu: X_0 \rightarrow Z/2Z$ will represent $1 \in H^0(G')$. Let $f: X_n \rightarrow Z/2Z$ represent $\alpha \in H^n(G')$ and for $i \geq 1$ let $f_i: X_i \rightarrow Z/2Z$ represent $\alpha_i \in H^i(G')$. The maps f, f_i may also be viewed as maps on the entire chain X by setting them equal to the zero map in all but the n -th, i -th component, respectively. By [3, Section 5.5], $\mathcal{A}(\gamma)$ is represented in $\text{Hom}_G(W \otimes X^l; Z/2Z)$ by $\mathcal{A}(\mu' + f + f_1 + \dots)$ where for any $g: X \rightarrow Z/2Z$

$$\mathcal{A}(g)(w_0 \otimes x_1 \otimes \dots \otimes x_l) = \varepsilon'(w_0)g(x_1) \dots g(x_l), \quad x_i \in X.$$

Since for $i \geq 1$ $f_i: X_j \rightarrow Z/2Z$ is zero for any $j \leq n$ it follows that if we wish to restrict $\mathcal{A}(\gamma)$ to $\text{Hom}_G((W \otimes X^l)_n; Z/2Z)$ we may use as a representative $\mathcal{A}(\mu' + f)$. By [3, first paragraph on p. 64], $\mathcal{A}(\mu' + f)$ represents $\text{cores}(\alpha)$ in $H^n(G)$. This completes the proof of the lemma.

Proof of Theorem 2. Let us apply diagram (5) to an element $q \in \hat{I}^n(E) \leq \hat{W}(E)$. By [6, Lemma 3.2], $w(q) = 1 + w_t(q) + \text{higher terms}$. By the lemma it follows that $\mathcal{A}(w(q)) = 1 + \text{cores}(w_t(q)) + \text{higher terms}$. We have $T(q) \in \hat{I}^n(F) \leq \hat{W}(F)$ and $w(T(q)) = 1 + w_t(T(q)) + \text{higher terms}$. Therefore it follows that $\text{cores}(w_t(q)) = w_t(T(q))$ as claimed.

Remarks. (1) If E/F is Galois of even degree then as in the main theorem of [8] we have $\text{cores} \circ w_t = w_t \circ T$ on $\hat{I}^n(E)$, modulo the kernel of $\text{res}: H^l(F) \rightarrow H^l(E)$.

(2) Whenever $h_F: k_\pi(F) \rightarrow H^n(F)$ is injective (also $h_F: k_\pi(E) \rightarrow H^n(E)$) we can

define $\iota : k_\pi(E) \rightarrow k_\pi(F)$. As previously noted h_F is known to be bijective for many cases. We have also shown (Theorem 1) that if we identify $k_\pi(E)$ as a subring of $H^\pi(E)$ via h_E then the Bass–Tate transfer ‘is’ corestriction. Theorem 2 then implies that $N \circ W_i = T \circ W_i$ on $\hat{I}^\pi(E)$, where N is the Bass–Tate transfer. It would be extremely enlightening to realize $\iota : k_\pi(E) \rightarrow k_\pi(F)$ from a purely K -theoretic perspective.

(3) By Tate’s result [10], if $T_2 : k_2(E) \rightarrow k_2(F)$ is the transfer map defined by Milnor in [1] then $\text{cores} \circ h_E = h_F \circ T_2$ on $k_2(E)$. If h_F is injective then it follows from Theorem 1 that $T_2 = N$, where N is the Bass–Tate transfer on $k_2(E)$. This is extremely interesting considering the divergence of their respective definitions.

References

- [1] H. Bass and J. Tate, The Milnor ring of a global field, *Algebraic K-theory II*, Lecture Notes in Math. No. 342 (Springer, Berlin–New York, 1973) 447–463.
- [2] A. Delzant, Definition des classes de Stiefel–Whitney d’un module quadratique, *C.R. Acad. Sci. Paris* 255 (1962) 1366–1368.
- [3] L. Evens, A generalization of the transfer map in the cohomology of groups, *Trans. Amer. Math. Soc.* 108 (1963) 54–65.
- [4] T.Y. Lam, *Algebraic Theory of Quadratic Forms* (Benjamin/Cummings, Reading, MA, 1973).
- [5] S. Lang, *Algebra* (Addison-Wesley, Reading, MA, 1965).
- [6] J. Milnor, Algebraic K -theory and quadratic forms, *Inventiones Math.* 9 (1970) 318–344.
- [7] J. Milnor, Introduction to Algebraic K -theory, *Annals of Math. Studies* No. 72 (Princeton University Press, 1971).
- [8] J.M. Shapiro, A Riemann–Roch type theorem for the Witt and Milnor rings of a field, *J. Pure Appl. Algebra* 15 (1979) 293–304.
- [9] S. Shatz, Profinite groups arithmetic and geometry, *Annals of Math. Studies* No. 67 (Princeton University Press, 1972).
- [10] J. Tate, Unpublished letter to S. Rosset (1977).
- [11] J. Tate, Relations between K_2 and Galois cohomology, *Inventiones Math.* 36 (1976) 257–274.